

# FuVar Notes

Naoki Pross – naoki.pross@ost.ch

Spring Semester 2021

## 1 Preface

These are just my personal notes of the FuVar course, and definitively not a rigorously constructed mathematical text. The good looking L<sup>A</sup>T<sub>E</sub>X typesetting may trick you into thinking it is rigorous, but really, it is not.

## 2 Derivatives of vector valued scalar functions

**Definition 1** (Partial derivative). A vector valued function  $f : \mathbb{R}^m \rightarrow \mathbb{R}$ , with  $\mathbf{v} \in \mathbb{R}^m$ , has a partial derivative with respect to  $v_i$  defined as

$$\partial_{v_i} f(\mathbf{v}) = f_{v_i}(\mathbf{v}) = \lim_{h \rightarrow 0} \frac{f(\mathbf{v} + h\mathbf{e}_i) - f(\mathbf{v})}{h}$$

**Proposition 1.** Under some generally satisfied conditions (continuity of  $n$ -th order partial derivatives) Schwarz's theorem states that it is possible to swap the order of differentiation.

$$\partial_x \partial_y f(x, y) = \partial_y \partial_x f(x, y)$$

**Definition 2** (Linearization). A function  $f : \mathbb{R}^m \rightarrow \mathbb{R}$  has a linearization  $g$  at  $\mathbf{x}_0$  given by

$$g(\mathbf{x}) = f(\mathbf{x}_0) + \sum_{i=1}^m \partial_{x_i} f(\mathbf{x}_0)(x_i - x_{i,0}),$$

if all partial derivatives are defined at  $\mathbf{x}_0$ .

**Theorem 1** (Propagation of uncertainty). Given a measurement of  $m$  values in a vector  $\mathbf{x} \in \mathbb{R}^m$  with values given in the form  $x_i = \bar{x}_i \pm \sigma_{x_i}$ , a linear approximation the error of a dependent variable  $y$  is computed with

$$y = \bar{y} \pm \sigma_y \approx f(\bar{\mathbf{x}}) \pm \sqrt{\sum_{i=1}^m (\partial_{x_i} f(\bar{\mathbf{x}}) \sigma_{x_i})^2}$$

**Definition 3** (Gradient vector). The *gradient* of a function  $f(\mathbf{x})$ ,  $\mathbf{x} \in \mathbb{R}^m$  is a vector containing the

derivatives in each direction.

$$\nabla f(\mathbf{x}) = \sum_{i=1}^m \partial_{x_i} f(\mathbf{x}) \mathbf{e}_i = \begin{pmatrix} \partial_{x_1} f(\mathbf{x}) \\ \vdots \\ \partial_{x_m} f(\mathbf{x}) \end{pmatrix}$$

**Definition 4** (Directional derivative). A function  $f(\mathbf{x})$  has a directional derivative in direction  $\mathbf{r}$  (with  $|\mathbf{r}| = 1$ ) given by

$$\frac{\partial f}{\partial \mathbf{r}} = \nabla_{\mathbf{r}} f = \mathbf{r} \cdot \nabla f$$

**Theorem 2.** The gradient vector always points towards the *direction of steepest ascent*.

## 3 Methods for maximization and minimization problems

**Method 1** (Find stationary points). Given a function  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}$ , to find its maxima and minima we shall consider the points

- that are on the boundary of the domain  $\partial D$ ,
- where the gradient  $\nabla f$  is not defined,
- that are stationary, i.e. where  $\nabla f = \mathbf{0}$ .

**Method 2** (Determine the type of stationary point for 2 dimensions). Given a scalar function of two variables  $f(x, y)$  and a stationary point  $\mathbf{x}_s$  (where  $\nabla f(\mathbf{x}_s) = \mathbf{0}$ ), we define the *discriminant*

$$\Delta = \partial_x^2 f \partial_y^2 f - \partial_y \partial_x^2 f$$

- if  $\Delta > 0$  then  $\mathbf{x}_s$  is an extrema, if  $\partial_x^2 f(\mathbf{x}_s) < 0$  it is a maximum, whereas if  $\partial_x^2 f(\mathbf{x}_s) > 0$  it is a minimum;
- if  $\Delta < 0$  then  $\mathbf{x}_s$  is a saddle point;
- if  $\Delta = 0$  we need to analyze further.

**Remark 1.** The previous method is obtained by studying the second directional derivative  $\nabla_{\mathbf{r}} \nabla_{\mathbf{r}} f$  at the stationary point in direction of a vector  $\mathbf{r} = \mathbf{e}_1 \cos(\alpha) + \mathbf{e}_2 \sin(\alpha)$

**Definition 5** (Hessian matrix). Given a function  $f : \mathbb{R}^m \rightarrow \mathbb{R}$ , the square matrix whose entry at the  $i$ -th row and  $j$ -th column is the second derivative of  $f$  first with respect to  $x_j$  and then to  $x_i$  is known as the *Hessian* matrix.  $(\mathbf{H}_f)_{i,j} = \partial_{x_i} \partial_{x_j} f$  or

$$\mathbf{H}_f = \begin{pmatrix} \partial_{x_1} \partial_{x_1} f & \cdots & \partial_{x_1} \partial_{x_m} f \\ \vdots & \ddots & \vdots \\ \partial_{x_m} \partial_{x_1} f & \cdots & \partial_{x_m} \partial_{x_m} f \end{pmatrix}$$

Because (almost always) the order of differentiation does not matter, it is a symmetric matrix.

**Method 3** (Determine the type of stationary point in higher dimensions). Given a scalar function of two variables  $f(x, y)$  and a stationary point  $\mathbf{x}_s$  (where  $\nabla f(\mathbf{x}_s) = \mathbf{0}$ ), we compute the Hessian matrix  $\mathbf{H}_f(\mathbf{x}_s)$ . Then we compute its eigenvalues  $\lambda_1, \dots, \lambda_m$  and

- if all  $\lambda_i > 0$ , the point is a minimum;
- if all  $\lambda_i < 0$ , the point is a maximum;
- if there are both positive and negative eigenvalues, it is a saddle point.

In the other cases, when there are  $\lambda_i \leq 0$  and/or  $\lambda_i \geq 0$  further analysis is required.

**Remark 2.** Recall that to compute the eigenvalues of a matrix, one must solve the equation  $(\mathbf{H} - \lambda \mathbf{I})\mathbf{x} = \mathbf{0}$ . Which can be done by solving the characteristic polynomial  $\det(\mathbf{H} - \lambda \mathbf{I}) = 0$  to obtain  $\dim(\mathbf{H})$   $\lambda_i$ , which when plugged back in result in an overdetermined system of equations.

**Method 4** (Quickly find the eigenvalues of a  $2 \times 2$  matrix). Let

$$m = \frac{1}{2} \operatorname{tr} \mathbf{H} = \frac{a+d}{2} \text{ and } p = \det \mathbf{H} = ad - bc,$$

then

$$\lambda = m \pm \sqrt{m^2 - p}.$$

**Method 5** (Search for a constrained extremum in 2 dimensions). Let  $n(x, y) = 0$  be a constraint in the search of the extrema of a function  $f : D \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$ . To find the extrema we look for points

- on the boundary  $\mathbf{u} \in \partial D$  where  $n(\mathbf{u}) = 0$ ;
- $\mathbf{u}$  where the gradient either does not exist or is  $\mathbf{0}$ , and satisfy  $n(\mathbf{u}) = 0$ ;
- that solve the system of equations

$$\begin{cases} \partial_x f(\mathbf{u}) \cdot \partial_y n(\mathbf{u}) = \partial_y f(\mathbf{u}) \cdot \partial_x n(\mathbf{u}) \\ n(\mathbf{u}) = 0 \end{cases}$$

**Method 6** (Search for a constrained extremum in higher dimensions, method of Lagrange multipliers). We wish to find the extrema of  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}$  under  $k < m$  constraints  $n_1 = 0, \dots, n_k = 0$ . To find the extrema we consider the following points:

- Points on the boundary  $\mathbf{u} \in \partial D$  that satisfy  $n_i(\mathbf{u}) = 0$  for all  $1 \leq i \leq k$ ,
- Points  $\mathbf{u} \in D$  where either
  - any of  $\nabla f, \nabla n_1, \dots, \nabla n_k$  do not exist, or
  - $\nabla n_1, \dots, \nabla n_k$  are linearly *dependent*,
 and that satisfy  $0 = n_1(\mathbf{u}) = \dots = n_k(\mathbf{u})$ .
- Points that solve the system of  $m+k$  equations

$$\begin{cases} \nabla f(\mathbf{u}) = \sum_{i=1}^k \lambda_i \nabla n_i(\mathbf{u}) & (m\text{-dimensional}) \\ n_i(\mathbf{u}) = 0 & \text{for } 1 \leq i \leq k \end{cases}$$

The  $\lambda$  values are known as *Lagrange multipliers*. The same calculation can be written more compactly by defining the  $m+k$  dimensional *Lagrangian*

$$\mathcal{L}(\mathbf{u}, \boldsymbol{\lambda}) = f(\mathbf{u}) - \sum_{i=1}^k \lambda_i n_i(\mathbf{u})$$

where  $\boldsymbol{\lambda} = \lambda_1, \dots, \lambda_k$  and then evaluating  $\nabla \mathcal{L}(\mathbf{u}, \boldsymbol{\lambda}) = \mathbf{0}$ .

## 4 Integration

**Remark 3.**

### License

This work is licensed under a “[CC BY-NC-SA 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/)” license.

